

Pro

LEBANON Scape®

FERTILIZER with TRIMEC® HERBICIDE BROADLEAF WEED CONTROL

20-0-7



METH-EX®

GUARANTEED ANALYSIS

Total Nitrogen (N)	20%
1.6% Water Insoluble Nitrogen*	
15.6% Urea Nitrogen	
2.8% Other Water Soluble Nitrogen*	
Soluble Potash (K ₂ O)	7%
Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.	
Chlorine (Cl) not more than	6.0% F699
*4.4% Slowly available Nitrogen from Methylene Ureas.	



0 88685 53597 0

22-53597

METH-EX and ProScape are registered trademarks of Lebanon Seaboard Corporation.

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

SUGGESTED SPREADER SETTINGS			
Spreader	4#/1,000 sq. ft.	Spreader	4#/1,000 sq. ft.
LebanonTurf	4½	Lesco (letter/Numeric)	J/21
Andersons AccuPro	M	PennMulch HVO	L
Earthway Rotary	16	ProScape SS	L
Gandy	27	Spyker	4½
Lely	6 II	Vicon (all models)	26

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, and other broadleaf weeds.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid.....	0.55%
(+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.12%
Dicamba: (3,6-dichloro-o-anisic acid)	0.05%

INERT INGREDIENTS

TOTAL

THIS PRODUCT CONTAINS:

11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
2.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.
1.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Contains the single isomer form of Mecoprop-p.

Isomer Specific by AOAC Methods.

Trimec® is a registered trademark of PBI/GORDON CORPORATION.

EPA Reg. No. 2217-581-961

(L) (D)
EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for Precautionary Statements.



Manufactured by:

Lebanon Seaboard Corporation
1600 E. Cumberland St. • Lebanon, PA 17042

NET WEIGHT 40 LBS. (18.14 KG)

FERTILIZER

with TRIMEC® HERBICIDE

BROADLEAF WEED CONTROL

20-0-7



METH-EX®

GUARANTEED ANALYSIS

Total Nitrogen (N) 20%
 1.6% Water Insoluble Nitrogen*
 15.6% Urea Nitrogen
 2.8% Other Water Soluble Nitrogen*
 Soluble Potash (K₂O) 7%

Derived From: Ammonium Sulfate, Methylene Ureas, Urea, Muriate of Potash.

Chlorine (Cl) not more than 6.0% F699
 *4.4% Slowly available Nitrogen from Methylene Ureas.



22-53597

METH-EX and ProScape are registered trademarks of Lebanon Seaboard Corporation.

Information regarding the contents and levels of metals in this product is available on the Internet at:
<http://www.regulatory-info-lebsea.com>

SUGGESTED SPREADER SETTINGS

Spreader	4#/1,000 sq. ft.	Spreader	4#/1,000 sq. ft.
LebanonTurf	4½	Lesco (letter/Numeric)	J/21
Andersons AccuPro	M	PennMulch HVO	L
Earthway Rotary	16	ProScape SS	L
Gandy	27	Spyker	4½
Lely	6 II	Vicon (all models)	26

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and evenness of terrain may require slightly different settings for desired coverage.

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, and other broadleaf weeds.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid.....	0.55%
(+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.12%
Dicamba: (3,6-dichloro-o-anisic acid)	0.05%

INERT INGREDIENTS 99.28%

TOTAL 100.00%

THIS PRODUCT CONTAINS:

11.0 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.

2.4 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.

1.0 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Contains the single isomer form of Mecoprop-p.

Isomer Specific by AOAC Methods.

Trimec® is a registered trademark of PBI/GORDON CORPORATION.

EPA Reg. No. 2217-581-961

(L) (D)
EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

KEEP OUT OF REACH OF CHILDREN CAUTION

See back panel for Precautionary Statements.



Manufactured by:

Lebanon Seaboard Corporation

1600 E. Cumberland St. • Lebanon, PA 17042

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS **HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

CAUTION: Harmful if absorbed through skin. Harmful if inhaled. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Avoid breathing dust.

When using this product, wear long-sleeved shirt, long pants, chemical-resistant gloves, socks and shoes.

User should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside, then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

FIRST AID

If in eyes:	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15 - 20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
If on skin or clothing:	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
If inhaled:	<ul style="list-style-type: none">• Move person to fresh air.• If person is not breathing, call 911 or ambulance, then give artificial respiration, preferable mouth-to-mouth, if possible.• Call a poison control center or doctor for further treatment advice.
If swallowed:	<ul style="list-style-type: none">• Call a poison control center or doctor immediately for further treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by a poison control center or doctor.• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also call toll free 1-877-800-5556 for emergency medical treatment information.	

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish and aquatic invertebrates and may adversely affect non-target plants. To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

Do not apply directly to or near water, storm drains, gutters, sewers, or drainage ditches. Do not apply within 25 feet of rivers, fish ponds, lakes, streams, reservoirs, marshes, estuaries, bays and oceans. Do not apply when windy. Apply this product directly to your lawn or garden, and sweep any product landing on the driveway, sidewalk, gutter, or street back onto the treated area. To prevent product run-off, do not over water the treated area(s) to the point of runoff or apply when raining or when rain is expected that day.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage and disposal.

PESTICIDE STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. **If empty:** Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

WHERE TO USE

This product is a Trimec® brand herbicide containing fertilizer, 2,4-D, mecoprop-p, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

This product is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example, hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses) and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings.

Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

This product may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

This product may be applied to newly established warm season turfgrass. Treatments should be delayed until 3 to 4 weeks after the sodding, springing, or plugging operations, and low application rates (½ of the recommendation for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Do not use this product on dichondra, carpetgrass, St. Augustinegrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

WEEDS CONTROLLED

Dandelion, chickweed, plantain, and clover are controlled effectively with this product. A partial list of susceptible weeds is shown below:

Bedstraw	Burdock	Clover	Ground Ivy	Knotweed	Peppergrass	Purslane	Shepherdspurse
Black Medic	Chicory	Dandelion	Healall	Lambsquarters	Pigweed	Ragweed	Speedwell
Buckhorn	Chickweed	Dock	Henbit	Morningglory	Plantain	Sheep Sorrel	Thistle

HOW MUCH TO APPLY

For Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, Bermudagrass, and Zoysiagrass -

Apply 139 to 218 pounds of this product per acre of established turfgrass. Or the equivalent application rate is 3.2 to 3.5 pounds of product per 1000 square feet of turfgrass.

Generally, the lower application rates within the specified range will provide satisfactory control of sensitive weed species. The higher the application rates within the specified range will be required for dense infestations of perennial weeds, and for adverse/extreme environmental conditions, or for weeds beyond the appropriate growth stages.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site.

The maximum number of broadcast applications per treatment site is 2 per year. Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds.

Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days. Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer.

Bag Size, Pounds	Desired Rate	Desired Rate	Treated Area For This Bag Size	
	Per Acre, Lbs	Per 1,000 Sq. Ft., Lbs.	Acres	Sq. Ft.
40	139-218	3.2 to 5	0.18 to 0.29	8,000 to 12,500

APPLICATION SUGGESTIONS

1. Spring and fall applications are recommended for the best control. However, this product can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 2 to 3 days before application.
3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

NOTE: Supplemental applications of lawn fertilizer may be needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

LIMITED WARRANTY AND DISCLAIMER

The manufacturer warrants only that the chemical composition of this product conforms to the ingredient statement given on the label, and that the product is reasonably suited for the labeled use when applied according to the Directions for Use.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE MANUFACTURER NEITHER MAKES NOR INTENDS ANY OTHER EXPRESS OR IMPLIED WARRANTIES, INCLUDING ANY WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE, WHICH ARE EXPRESSLY DISCLAIMED. This limited warranty does not extend to the use of the product inconsistent with label instructions, warnings or cautions, or to use of the product under abnormal conditions such as drought, excessive rainfall, tornadoes, hurricanes, etc. These factors are beyond the control of the manufacturer or the seller. Any damages arising from a breach of the manufacturer's warranty shall be limited to direct damages, and shall not include indirect or consequential damages such as loss of profits or values, except as otherwise provided by law.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, the terms of this Limited Warranty and Disclaimer cannot be varied by any written or verbal statements or agreements. No employee or agent of the seller is authorized to vary or exceed the terms of this Limited Warranty and Disclaimer in any manner.