

COVERS UP TO **15,600** SQ. FT.

**ProScape** LEBANON

**FERTILIZER WITH**

**TRIMEC\* HERBICIDE**  
BROADLEAF WEED CONTROL

# ProScape<sup>®</sup> **LEBANON**

## HOMOGENEOUS FERTILIZER

WITH

## TRIMEC<sup>®</sup> HERBICIDE BROADLEAF WEED CONTROL

### 20-4-10



**GUARANTEED ANALYSIS**

Total Nitrogen (N)	20%
11.0% Ammoniacal Nitrogen	
4.8% Water Insoluble Nitrogen	
1.2% Urea Nitrogen	
3.0% Other Water Soluble Nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> )	4%
Soluble Potash (K <sub>2</sub> O)	10%
Total Iron (Fe)	0.5%
0.05% Water Soluble Iron (Fe)	
Nutrient Sources: Ammonium Phosphate, Ammonium Sulfate, Isobutylidene Diurea, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.	
Chlorine (Cl) not more than	9.0%
*3.0% Slowly Available Nitrogen from Methylene Ureas.	F699

54473

Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, and other broadleaf weeds.

**ACTIVE INGREDIENTS:**

2,4-dichlorophenoxyacetic acid	0.69%
(+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid	0.15%
Dicamba: 3,6-dichloro-o-anisic acid	0.06%

<b>INERT INGREDIENTS</b>	<b>99.10%</b>
<b>TOTAL</b>	<b>100.00%</b>

**THIS PRODUCT CONTAINS:**

13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.  
3.0 lbs. (+)-(R)-2-(2-methyl-4-chlorophenoxy)propionic acid equivalent per ton.  
1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

EPA Reg. No. 2217-795-961

TRIMEC<sup>®</sup> is a registered trademark of PBI/GORDON CORPORATION.

(L)

(D)

EPA Est. 961-PH-01-961-IL-01

Superscript used is first letter of lot number.

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION**

See Back Panel for Precautionary Statements.

Distributed by

**Lebanon Seaboard Corporation**

1600 E. Cumberland St. • Lebanon, PA 17042

[www.LebanonTurf.com](http://www.LebanonTurf.com)

NET WEIGHT **50** LBS. (22.68 KG)

**TRIMEC\* HERBICIDE**  
BROADLEAF WEED CONTROL

**FERTILIZER WITH**

**ProScape** LEBANON

SPECIMEN LABEL

# 20-4-10



### GUARANTEED ANALYSIS

Total Nitrogen (N) . . . . .	20%
11.0% Ammoniacal Nitrogen	
4.8% Water Insoluble Nitrogen	
1.2% Urea Nitrogen	
3.0% Other Water Soluble Nitrogen*	
Available Phosphate (P <sub>2</sub> O <sub>5</sub> ) . . . . .	4%
Soluble Potash (K <sub>2</sub> O) . . . . .	10%
Total Iron (Fe) . . . . .	0.5%
0.05% Water Soluble Iron (Fe)	

Nutrient Sources: Ammonium Phosphate, Ammonium Sulfate, Isobutylidene Diurea, Methylene Ureas, Urea, Muriate of Potash, Ferric Oxide, Ferrous Sulfate.

Chlorine (Cl) not more than . . . . .	9.0%
*3.0% Slowly Available Nitrogen from Methylene Ureas.	F699

NOTICE: This product contains the secondary nutrient iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly after application by sweeping or blowing. Do not wash off with water.

**54473**

**Controls Dandelion, Chickweed, Knotweed, Plantain, Henbit, and other broadleaf weeds.**

### ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid . . . . .	0.69%
(+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid . . . . .	0.15%
Dicamba: 3,6-dichloro-o-anisic acid . . . . .	0.06%

<b>INERT INGREDIENTS</b> . . . . .	<b>99.10%</b>
<b>TOTAL</b> . . . . .	<b>100.00%</b>

### THIS PRODUCT CONTAINS:

- 13.8 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
- 3.0 lbs. (+)-(R)-2-(2 methyl-4-chlorophenoxy)propionic acid equivalent per ton.
- 1.2 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.

Isomer Specific by AOAC Methods.

Contains the single isomer form of Mecoprop-p.

EPA Reg. No. 2217-795-961

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

(L) (D)

EPA Est. 961-PA-01:961-IL-01

Superscript used is first letter of lot number.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

See Back Panel for Precautionary Statements.

Distributed by

**Lebanon Seaboard Corporation**

1600 E. Cumberland St. • Lebanon, PA 17042

[www.LebanonTurf.com](http://www.LebanonTurf.com)

SPECIMEN LABEL

# LEBANON **ProScape**®

## HOMOGENEOUS FERTILIZER WITH **TRIMEC**® HERBICIDE BROADLEAF WEED CONTROL

**STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.**

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS **CAUTION**

Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes. After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hands and exposed skin with soap and water.

#### FIRST AID

- IF IN EYES:**
- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
  - Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
  - Call a poison control center or doctor for treatment advice.
- IF ON SKIN  
OR CLOTHING:**
- Take off contaminated clothing.
  - Rinse skin immediately with plenty of water for 15-20 minutes.
  - Call a poison control center or doctor for treatment advice.
- IF SWALLOWED:**
- Call a poison control center or doctor immediately for treatment advice.
  - Have person sip a glass of water if able to swallow.
  - Do not induce vomiting unless told to do so by a poison control center or doctor.
  - Do not give anything by mouth to an unconscious person.
- IF INHALED:**
- Move person to fresh air.
  - If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.
  - Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment. For Emergency Medical Information call (888) 208-1368. You may also call toll-free, 1-877-800-5556 for emergency medical treatment information.

#### ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. For terrestrial uses, do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment washwater. When cleaning equipment, do not pour the washwater on the ground; spray or drain over a large area away from wells and other water sources. Do not apply when weather conditions favor drift from target area.

Most cases of groundwater contamination involving phenoxy herbicides such as MCP and 2,4-D have been associated with mixing/loading and disposal sites. Caution should be exercised when handling MCP and 2,4-D pesticides at such sites to prevent contamination of groundwater supplies. Use of closed systems for mixing and transferring this pesticide will reduce the probability of spills. Placement of the mixing/loading equipment on an impervious pad to contain spills will help prevent groundwater contamination.

#### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.

#### STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals.

**DISPOSAL:**

**If empty** - Do not reuse this container. Place in trash or offer for recycling if available.

**If partially filled** - Call your local solid waste agency or toll free number (1-800-CLEANUP) for disposal instructions. Never place unused product down any indoor or outdoor drain.

#### WHERE TO USE

This product is a Trimec® brand herbicide containing fertilizer, 2,4-D, mecoprop, and dicamba. This granular weed and feed provides plant nutrients and controls a broad spectrum of broadleaf weeds in turfgrass. Supplemental nitrogen, phosphorus, and potash should be applied according to soil tests or State Extension Service recommendations.

SPECIMEN LABEL

This product is intended to be applied by commercial applicators, noncommercial users, lawn care operators, and landscape personnel. Cool season and warm season grasses established in institutional, ornamental, and residential sites may be treated.

Institutional sites are defined as turf areas around properties or facilities providing a service to public or private organizations; for example areas around hospitals, nursing homes, schools, museums, libraries, sport facilities (including golf courses), and office buildings may be treated. Ornamental sites include turfgrass established around residences, parks, streets, retail outlets, cemeteries, and institutional buildings. Finally, residential sites are defined as areas associated with the household or home life including, but not limited to apartment complexes, condominiums, and patient care areas of nursing homes, mental institutions, hospitals, or convalescent homes.

This product may be applied to newly seeded cool season turfgrass. Applications are recommended only after the third mowing or after the grass seedlings have tillered and have developed an adequate secondary root system.

This product may be applied to newly established warm season turfgrass. Treatments should be delayed until 3-4 weeks after the sodding, sprigging, or plugging operations, and low application rates (1/2 of the recommendation for established turfgrass) are suggested.

A uniform broadcast distribution of product on the treated area is essential. Do not exceed the specified application rate for any area and use care in the application of this product within the dripline of trees, shrubs, and other ornamentals.

Do not use this product on dichondra, carpetgrass, St. Augustinegrass, closely-mowed bentgrass putting greens, or turf containing desirable clovers.

## WEEDS CONTROLLED

Dandelion, chickweed, plantain, and clover are controlled effectively with this product. A partial list of susceptible weeds is shown below:

Bedstraw	Burdock	Clover	Ground ivy	Knotweed	Peppergrass	Purslane	Shepherdspurge
Black medic	Chicory	Dandelion	Healall	Lambsquarters	Pigweed	Ragweed	Speedwell
Buckhorn	Chickweed	Dock	Henbit	Morningglory	Plantain	Sheep sorrel	Thistle

## HOW MUCH TO APPLY

**For Kentucky Bluegrass, Fescue spp., Perennial Ryegrass, Bermudagrass, and Zoysiagrass:** Apply 140.5 to 174.2 pounds of this product per acre of established turfgrass. Or the equivalent application rates are 3.2 to 4.0 pounds of product per 1,000 square feet of turfgrass.

The maximum application rate to turf is 1.2 pounds 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

Sequential or follow-up applications may be required for dense infestations of perennial and biennial weeds. Also, adverse environmental conditions prior or after application may reduce the effectiveness. If required, the treatment interval between the first and second applications should be approximately 30 days.

Drop spreaders, rotary spreaders, and other granular equipment should be calibrated for accurate and uniform applications of this product. Calibrate the equipment by applying the product over a measured or known area and adjust the equipment according to the recommendations of the equipment manufacturer.

Bag Size, Lbs.	Desired Rate	Desired Rate	Treated Area For This Bag Size	
	Per Acre, Lbs.	Per 1,000 Sq. Ft., Lbs.	Acres	Sq. Ft.
50	140.5	3.2	0.36	15,600
	158.4	3.6	0.32	13,900
	174.2	4.0	0.29	12,500

## APPLICATION SUGGESTIONS

1. Spring and fall applications are recommended for the best control. However, this product can be applied any time broadleaf weeds are present and actively growing.
2. Mow the turfgrass 2 to 3 days before application.
3. Water the turfgrass thoroughly (>1/2 inch) at least 1 to 2 days before application.
4. Schedule applications in the morning when dew is present. Or, sprinkle or irrigate lightly (<1/4 inch) before application to provide maximum control.
5. Rainfall or irrigation within 24 hours after application may reduce the effectiveness. Do not irrigate or water the turfgrass within 24 hours after application.

NOTE: Supplemental applications of lawn fertilizer may be needed to achieve a minimum 1 pound nitrogen per 1,000 sq. ft.

## SUGGESTED SPREADER SETTINGS

Spreader	Settings		Spreader	Settings	
	3.2#/1000 Sq. Ft.	4.0#/1000 Sq. Ft.		3.2#/1000 Sq. Ft.	4.0#/1000 Sq. Ft.
Cyclone	4	5	PrizeLAWN® BF & CBR III	H	I
Earthway (Rotary)	13	15	PrizeLAWN® CBR II	3 1/2	4
Ezee Flow (109)	C2	C3	PrizeLAWN® LF-1	2 1/2	3
Gandy	30	32	Republic Rotary	4	5 1/2
Lely (HR,1250,1500,2010)	5II	6II	Scotts AccuPro	H	M
Lely (W)	5II	6II	Scotts (Rotary) R7X	F	H
Lesco (letter dial)	H	J	Scotts (Rotary) R8A	J	K
Lesco (numeric dial)	16	21	Spyker	2	4
Ortho Rotary	4	5 1/2	Vicon (all models)	20	28

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

## WARRANTY

The Distributor warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to Distributor, and Buyer assumes the risk of any such use.

DISTRIBUTED BY:  
**LEBANON SEABOARD CORPORATION**  
 1600 EAST CUMBERLAND STREET  
 LEBANON, PA 17042  
[www.LebanonTurf.com](http://www.LebanonTurf.com)

SPECIMEN LABEL