

NET WT. 20 LBS. (9.08 kg)
COVERS UP TO 10,000 SQ. FT.

35799



LEBANON®

BROADLEAF WEED KILLER

with **TRIMEC**® HERBICIDE

BROADLEAF WEED KILLER
with **TRIMEC**® HERBICIDE

CONTROLS: Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds, including those listed on the back panel.

ACTIVE INGREDIENTS:	
2,4-dichlorophenoxyacetic acid	1.3677%
2-(2-methyl-4-chlorophenoxy) propionic acid	0.6120%
Dicamba: 3,6-dichloro-o-anisic acid	0.1290%
INERT INGREDIENTS:	97.8913%
TOTAL	100.0000%
THIS PRODUCT CONTAINS:	
27.35 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.	
12.24 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.	
2.58 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.	
Isomer Specific by AOAC Methods	

EPA REG. NO. 2217-640-961 (L) (D)
EPA Est. 961-PA-01:961-JL-01
Superscript used is first letter of lot number.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water.
For eyes, flush with water for 15 minutes and get medical attention.
See back panel for additional Precautionary Statements.



DISTRIBUTED BY
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.lebsea.com

35799



CONTROLS: Dandelion, Chickweed, Knotweed, Plantains, Henbit, Spurge and many other broadleaf weeds, including those listed on the back panel.

ACTIVE INGREDIENTS:

2,4-dichlorophenoxyacetic acid	1.3677%
2-(2-methyl-4-chlorophenoxy) propionic acid	0.6120%
Dicamba: 3,6-dichloro-o-anisic acid	0.1290%

INERT INGREDIENTS:	97.8913%
TOTAL	100.0000%

THIS PRODUCT CONTAINS:

27.35 lbs. 2,4-dichlorophenoxyacetic acid equivalent per ton.
12.24 lbs. 2-(2-methyl-4-chlorophenoxy) propionic acid equivalent per ton.
2.58 lbs. 3,6-dichloro-o-anisic acid equivalent per ton.
Isomer Specific by AOAC Methods

EPA REG. NO. 2217-640-961

(L) (D)
EPA Est. 961-PA-01:961-IL-01
Superscript used is first letter of lot number.

TRIMEC® is a registered trademark of PBI/GORDON CORPORATION.

KEEP OUT OF REACH OF CHILDREN


CAUTION

STATEMENT OF PRACTICAL TREATMENT

In case of contact, wash skin with plenty of soap and water.
For eyes, flush with water for 15 minutes and get medical attention.
See back panel for additional Precautionary Statements.



DISTRIBUTED BY
LEBANON SEABOARD CORPORATION
1600 EAST CUMBERLAND STREET
LEBANON, PA 17042
www.lebsea.com



NET WT. 20 LBS. (9.08 kg)
COVERS UP TO 10,000 SQ. FT.



BROADLEAF WEED KILLER

with **TRIMEC[®] HERBICIDE**

STOP! READ THE ENTIRE LABEL FIRST. OBSERVE ALL PRECAUTIONS AND FOLLOW DIRECTIONS CAREFULLY.

PRECAUTIONARY STATEMENTS

Hazards To Humans and Domestic Animals

CAUTION

Harmful if swallowed. Do not inhale dust. Do not allow contact with skin, eyes or clothing. When using this product, wear long-sleeved shirt, long pants, socks and shoes.

After using this product, remove clothing and launder separately before reuse, and promptly and thoroughly wash hand and exposed skin with soap and water.

ENVIRONMENTAL HAZARDS

Drift or runoff may adversely affect nontarget plants. Do not apply directly to water. Do not contaminate water when disposing of equipment washwater. Do not contaminate domestic or irrigation waters. Do not apply when weather conditions favor drift from target area.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not allow people (other than applicator) or pets on treatment area during application. Do not enter treatment area until spray has dried or dust has settled.

STORAGE AND DISPOSAL

STORAGE: Store in original container and keep any bags containing unused product in a locked storage area inaccessible to children and domestic animals. Do not store near food or feedstuffs.

PESTICIDE DISPOSAL: Securely wrap bag in several layers of newspaper and discard in trash.

CONTAINER DISPOSAL: Do not reuse bag. Discard bag in trash.

USE PRECAUTIONS

Do not allow the material to drift onto flowers, vegetables or ornamental shrubbery. Apply when the air is calm. Do not apply to newly seeded lawns. Do not apply in areas underlaid by roots of desirable trees and shrubs. Do not exceed specified dosages for any area; be particularly careful within dripline of tree and other ornamental species. During application, close spreader openings when turning. Do not contaminate domestic or irrigation waters.

INSTRUCTIONS: TRIMEC[®] controls a wide range of lawn weeds including dandelion, chickweed, knotweed, plantain, henbit, spurge and many others commonly found in home lawns.

Possible retreatment may be necessary if (a) lawn is heavily overrun by weeds, or, (b) if adverse low moisture conditions prevail and weeds are in a state of poor growth. A second application may be made in approximately 30 days.

Established turf of warm season grasses includes bermudagrass, centipedegrass, and zoysiagrass. Established turf of cool season grasses includes Kentucky bluegrass, perennial ryegrass, tall fescue, and turf-type tall fescues. Do not apply this product on carpetgrass, dichondra, St. Augustinegrass, or established turf containing desirable clovers.

The maximum application rate to turf is 1.2 lbs. 2,4-D acid equivalent per acre per application per site. The maximum number of broadcast applications per treatment site is 2 per year.

1. Mow lawn to normal height 1 to 2 days before application.
2. Water lawn thoroughly at least 1 day before application to sustain moisture until the next watering (see step 4).
3. Apply when weeds are young and actively growing, preferably in the morning when dew is on the grass. If grass is not moist at time of application, sprinkle lightly with water to hold the material and prevent dust drift onto nontarget plants.
4. Do not wash from weed leaves for 1 to 2 days after application. At this time a thorough watering should be made.

SPREADER SETTINGS			
Spreader	Setting	Spreader	Setting
Scotts R8A	J	Scotts (Rotary)	4 1/2
Earthway (Rotary)	16	Ortho (Drop)	8
Lesco	E	Ortho (Rotary)	3
Spyker	4	Jet-Spred® II	4
Cyclone	4	Drop-Spred® III	7
Scotts (Drop)	6	Jet-Spred® III	4 1/2

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage. The above spreader settings are approximate. Calibrate your spreader on 100 square feet and adjust to apply at the rate of 1 lb. of product per 500 square feet.

WEEDS

Bedstraw	Dandelion	Lespedeza	Poison Ivy	Spurge
Black Medic	Dock	Mallow	Poison Oak	Thistle
Buckhorn	Ground Ivy	Morningglory	Purslane	Wild Carrot
Burdock	Healall	Oxalis	Ragweed	Wild Garlic
Chicory	Henbit	Peppergrass	Sheep Sorrel	Wild Lettuce
Chickweed	Knotweed	Pigweed	Shepherdspurse	Wild Onion
Clover	Lambsquarters	Plantain	Speedwell	Yarrow

WARRANTY

The Distributor warrants that this product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, expressed or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to Distributor, and Buyer assumes the risk of any such use.

DISTRIBUTED BY
LEBANON SEABOARD CORPORATION
 1600 EAST CUMBERLAND STREET
 LEBANON, PA 17042
www.lebsea.com

35799