

NET WT. 40 LBS. (18.15 KG)  
COVERS UP TO 22,800 SQ. FT.

# Lebanon TEAM\* 2G HERBICIDE



Lebanon TEAM\* 2G  
HERBICIDE

Lebanon TEAM\* 2G  
HERBICIDE

SPECIMEN LABEL

TEAM<sup>®</sup> is a selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

Bahiagrass	Centipede	St. Augustine
Bermudagrass	Fescue	Zoysiagrass
Bluegrass (perennial)	Ryegrass (perennial)	Bentgrass fairways and other high cut Bentgrass areas

**ACTIVE INGREDIENTS:**

benefin: N-butyl-N-ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-p-toluidine	1.33%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro-N,N-dipropyl-p-toluidine	0.67%

<b>INERT INGREDIENTS</b>	98.00%
<b>TOTAL</b>	100.00%

Contains 1 pound active ingredient per 50 pound bag.  
EPA Reg. No. 62719-137-961 (L) (D)  
EPA Est. 961-PA-01961-L-01  
\*Trademark of Dow AgroSciences LLC. Superscript used is first letter of lot number.

**KEEP OUT OF REACH OF CHILDREN  
CAUTION PRECAUCION**

Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

Refer to back panel for additional precautionary information and Directions for Use.  
Notice: Read the entire label. Use only according to label directions. Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of this label. If terms are unacceptable, return at once unopened.  
In case of emergency endangering the environment involving this product, call 1-800-424-9300.  
Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**PRECAUTIONARY STATEMENTS  
HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
CAUTION PRECAUCION**

Precaucion al usuario: Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.  
Causes Eye Irritation • Harmful If Swallowed, Inhaled, Or Absorbed Through The Skin.

Avoid breathing dust and contact with skin, eyes, or clothing. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

**FIRST AID**

**IF SWALLOWED:** Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.  
**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.  
**IF IN EYES:** Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.  
**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.  
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-208-1388 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to fish. Do not apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

MANUFACTURED BY  
**LEBANON SEABOARD CORPORATION**  
1600 EAST CUMBERLAND STREET  
LEBANON, PA 17042  
www.LebanonTurf.com

TEAM\* is a selective preemergence herbicide for the control of crabgrass and most other annual weed grasses on lawns and golf courses in established:

Bahiagrass	Centipedegrass	St. Augustinegrass
Bermudagrass	Fescue	Zoysiagrass
Bluegrass (perennial)	Ryegrass (perennial)	Bentgrass fairways and other high cut Bentgrass areas

**ACTIVE INGREDIENTS:**

benefin: <i>N</i> -butyl- <i>N</i> -ethyl- $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>p</i> -toluidine	1.33%
trifluralin: $\alpha,\alpha,\alpha$ -trifluoro-2,6-dinitro- <i>N,N</i> -dipropyl- <i>p</i> -toluidine	0.67%

**INERT INGREDIENTS** ..... 98.00%

**TOTAL** ..... 100.00%

Contains 1 pound active ingredient per 50 pound bag.  
 EPA Reg. No. 62719-137-961  
 \*Trademark of Dow AgroSciences LLC.

(L) (D)  
 EPA Est. 961-PA-01:961-IL-01  
 Superscript used is first letter of lot number.

**KEEP OUT OF REACH OF CHILDREN  
 CAUTION PRECAUCION**

**Precaucion al usuario:** Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

**Refer to back panel for additional precautionary information and Directions for Use.**

**Notice:** Read the entire label. Use only according to label directions. **Before using this product, read Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies at end of this label. If terms are unacceptable, return at once unopened.**

In case of emergency endangering the environment involving this product, call 1-800-424-9300.

Agricultural Chemical: Do not ship or store with food, feeds, drugs or clothing.

**PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

**CAUTION PRECAUCION**

**Precaucion al usuario:** Si usted no lee inglés, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.  
**Causes Eye Irritation • Harmful If Swallowed, Inhaled, Or Absorbed Through The Skin.**

**Avoid breathing dust and contact with skin, eyes, or clothing. Use eye protection and protective clothing such as coveralls, a long-sleeved shirt, and impermeable gloves when handling this product. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.**

**FIRST AID**

**IF SWALLOWED:** Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call poison control center or doctor for treatment advice.

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-888-208-1368 for emergency medical treatment information.

**ENVIRONMENTAL HAZARDS**


This pesticide is toxic to fish. Do not apply directly to water, areas where surface water is present, or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

**MANUFACTURED BY  
 LEBANON SEABOARD CORPORATION**

1600 EAST CUMBERLAND STREET

LEBANON, PA 17042

[www.LebanonTurf.com](http://www.LebanonTurf.com)



# Lebanon TEAM\* 2G HERBICIDE



## DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Read all Directions for Use carefully before applying.

**Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purposes.**

## STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

**STORAGE:** Store in original container only. In case of leak or spill, contain material and dispose as waste.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

**CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

## GENERAL INFORMATION

This product is a preemergence herbicide which is applied to established turfgrasses for the control of annual bluegrass (*Poa annua*), smooth and hairy crabgrass, goosegrass (silver crabgrass or crowfoot), barnyardgrass (watergrass), and green and yellow foxtail. This product also provides partial control of prostrate spurge (*Euphorbia supina*), spotted spurge (*Euphorbia maculata*), creeping spurge (*Euphorbia serpens*), creeping woodsorrel (*Oxalis corniculata*), and yellow woodsorrel (*Oxalis stricta*). This product controls susceptible annual weeds by killing their seeds as they germinate. Apply this product before these annual weeds begin to germinate since it does not control established weeds.

When this product is applied as directed, the following established turfgrasses are tolerant:

Bahiagrass	Centipedegrass	Kentucky bluegrass	St. Augustinegrass	Bentgrass Fairways and other
Bermudagrass	Fescue	Perennial ryegrass	Zoysiagrass	high cut Bentgrass areas

This product will not harm most nearby established ornamental trees and shrubs.

Do not apply this product to sod farms, turfgrasses grown for seed, farms, greenhouses or nurseries, including plants or plant materials grown for sale or research purposes.

## GENERAL USE PRECAUTIONS

Apply this product to the turf prior to the germination of weed grasses since it will not control established weeds. It may thin established annual bluegrass (*Poa annua*) turfs and fine-leaved fescues at rates above 100 lbs. per acre (2 lbs. active ingredient). This product should not be applied in the spring to turfgrass planted the previous fall. In bermudagrass areas that have been over-seeded with winter grasses, a spring application of this product may thin the over-seeded grasses. Because of varying cultural practices, this product should not be used on putting greens. It should not be used on dichondra. It should not be applied to newly sprigged areas of bermudagrass, St. Augustinegrass, centipedegrass, or zoysiagrass until these turfs are well established.

## APPLICATION DIRECTIONS

### Rates, Frequency, and Timing of Application

Successful preemergence control of the annual grass weeds listed on this label requires proper timing of application.

Apply this product approximately 1 to 2 weeks prior to the germination of annual grass weeds. All summer annual grasses do not germinate at the same time. If application timing does not coincide with the normal germination period of any of the annual grass weeds listed on this label, weed control results may be erratic or poor. Optimum weed control performance should be obtained if treated areas are irrigated soon after application.

Any cultural practices that disturb the soil, such as aeration or verticutting, should be done prior to application of this product.

### 1. Spring Germinating Summer Annual Grasses - Crabgrasses, Goosegrasses, Foxtails, and Barnyardgrass

#### A. Northern (Cool Season) Turfgrasses

For preemergence control, apply 75 to 150 lbs. per acre (1.5 to 3 lbs. active ingredient) of this product in the spring 1 to 2 weeks prior to the onset of conditions favorable for annual weed grass seed germination. Where heavy goosegrass pressure exists, use the rate of 150 lbs. per acre of this product. If split applications are desired, apply a cumulative rate of 150 lbs. per acre (3 lbs. active ingredient) and make the second application 8 to 10 weeks after the first.

#### B. Southern (Warm Season) Turfgrasses

For preemergence control, apply 100 to 150 lbs. per acre (2 to 3 lbs. active ingredient) of this product in the late winter or early spring 1 to 2 weeks prior to the onset of conditions favorable for annual weed grass seed germination.

In areas of the United States where germination of the annual weed grass species continues for extended periods of time, a second application of this product at the recommended rate may be made 10 to 12 weeks after initial treatment for continued control.

### 2. *Poa annua* (Annual Bluegrass)

In areas of heavy infestation of annual bluegrass, its elimination will temporarily result in thin turfgrass areas. Proper fertilization, irrigation, and soil incorporated reseeding will encourage existing desirable turfgrasses and newly planted seedling turfgrasses to fill in thin areas previously occupied by annual bluegrass.

#### A. Northern (Cool Season) Turfgrasses

For preemergence control, apply 75 to 100 lbs. per acre (1.5 to 2 lbs. active ingredient) of this product in late summer or early fall and again in late winter or spring 1 to 2 weeks prior to the expected germination of annual bluegrass.

#### B. Southern (Warm Season) Turfgrasses

For preemergence control, apply 150 lbs. per acre (3 lbs. active ingredient) of this product in the late summer or early fall 1 to 2 weeks prior to the expected germination of annual bluegrass. In southern areas of the United States where germination of annual bluegrass continues for extended periods of time, a second application of this product at 150 lbs. per acre can be made 10 to 12 weeks after the initial treatment.

For application to smaller areas, apply the following amounts of TEAM 2G:

<u>TEAM 2G per Acre</u>	<u>TEAM 2G per 1,000 Sq. Ft.</u>
75 Lbs.	1 <sup>3</sup> / <sub>4</sub> Lbs.
100 Lbs.	2 <sup>1</sup> / <sub>4</sub> Lbs.
150 Lbs.	3 <sup>1</sup> / <sub>2</sub> Lbs.

## RESEEDING

Chemicals that effectively control annual weed seeds may also affect new desirable turfgrass seedlings. When reseeding, it is essential that proper cultural practices such as soil cultivation, irrigation, and fertilization be followed. For satisfactory reseeding results after using this product, the amount of seed used should be increased and equipment designed to place seed in contact with the soil should be employed. Reseeding should be delayed 8 weeks after application of this product at the 75 to 100 lbs. per acre (1.5 to 2 lbs. active ingredient) and 12 to 16 weeks at the 100 to 150 lbs. per acre (2 to 3 lbs. active ingredient).

## Application Equipment

Apply this product evenly over the turfgrass area without skips or misses. A more uniform application can be made by spreading half the required amount over the area and then applying the remaining half at right angles. Both drop type and rotary type spreaders may be used. Avoid the use of spreaders which would apply the material in narrow, concentrated bands. For best results, use equipment designed to apply granular herbicides and insecticides. Calibrate applicators prior to use according to the manufacturer's directions. Check frequently to be sure equipment is working properly and distributing granules uniformly.

SPECIMEN LABEL

### Helpful Application Techniques for Applying TEAM 2G

1. When using a drop type spreader, a splash board mounted under the hopper will help dispense granules more evenly. A chain fastened to the side of your spreader will leave a marked or scuffed turf area to follow and help prevent missed areas.
2. When using tractor mounted or pull-type rotary applicators to apply this product, it is suggested that early morning applications be made when the wind is calm and dew is on the ground. Wheel tracks are easy to follow and granules can be spread uniformly.
3. Small area application is easy using a small rotary applicator. To insure proper coverage and avoid missed areas, early morning treatment when wheel tracks are evident in the dew and the wind is calm are ideal.

#### Appropriate Spreader Settings As A Guide for Calibration (For applying TEAM 2G in one pass over the turf.)

**NOTE:** These settings are provided as *suggested* starting points in calibrating each individual spreader. Since no two spreaders are alike, these settings are not intended to be used as absolute recommendations by the Seller or the spreader manufacturer.

SUGGESTED SPREADER SETTINGS							
Spreader	Settings			Spreader	Settings		
	1 <sup>3</sup> / <sub>4</sub> #/1000 Sq. Ft.	2 <sup>1</sup> / <sub>4</sub> #/1000 Sq. Ft.	3 <sup>1</sup> / <sub>2</sub> #/1000 Sq. Ft.		1 <sup>3</sup> / <sub>4</sub> #/1000 Sq. Ft.	2 <sup>1</sup> / <sub>4</sub> #/1000 Sq. Ft.	3 <sup>1</sup> / <sub>2</sub> #/1000 Sq. Ft.
Cyclone	3 <sup>1</sup> / <sub>2</sub>	4	4 <sup>1</sup> / <sub>2</sub>	Prizelawn BF & CBR III	E	J	L
Earthway Rotary	14	15	17	Prizelawn CBR II	2 <sup>1</sup> / <sub>2</sub>	4	4 <sup>1</sup> / <sub>2</sub>
Ezee Flow (109)	C2	C2 <sup>1</sup> / <sub>2</sub>	C3	Prizelawn LF-1	2	2 <sup>1</sup> / <sub>2</sub>	3 <sup>1</sup> / <sub>2</sub>
Gandy	25	29	32	Republic Rotary	2 <sup>1</sup> / <sub>2</sub>	3	4 <sup>1</sup> / <sub>2</sub>
Lely (HR,1250,1500,2010)	5 <sup>1</sup> / <sub>2</sub> II	6 <sup>1</sup> / <sub>2</sub> II	7II	Scotts AccuPro	F	H	K
Lely (W)	5 <sup>1</sup> / <sub>2</sub> II	6 <sup>1</sup> / <sub>2</sub> II	7II	Scotts (Rotary) R7X	E	G	H
Lesco (letter dial)	D	G	I	Scotts (Rotary) R8A	J	K	L
Lesco (numeric dial)	10	15	17	Spyker	1 <sup>1</sup> / <sub>2</sub>	2	3
Ortho Rotary	2 <sup>1</sup> / <sub>2</sub>	3	4 <sup>1</sup> / <sub>2</sub>	Vicon (all models)	18	24	26

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation, and evenness of terrain may require slightly different settings for desired coverage.

### TERMS AND CONDITIONS OF USE

If terms of the following Warranty Disclaimer, Inherent Risks of Use, and Limitation of Remedies are not acceptable, return unopened package at once to the seller for a full refund of purchase price paid. Otherwise, use by the buyer or any other user constitutes acceptance of the terms under Warranty Disclaimer, Inherent Risks of Use and Limitations of Remedies.

#### WARRANTY DISCLAIMER

Seller warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions, subject to the inherent risks set forth below. SELLER MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

#### INHERENT RISKS OF USE

It is impossible to eliminate all risks associated with use of this product. Plant injury, lack of performance, or other unintended consequences may result because of such factors as use of the product contrary to label instructions (including conditions noted on the label, such as unfavorable temperature, soil conditions, etc.), abnormal conditions (such as excessive rainfall, drought, tornadoes, hurricanes), presence of other materials, the manner of application, or other factors, all of which are beyond the control of the Seller. All such risks shall be assumed by Buyer.

#### LIMITATION OF REMEDIES

The exclusive remedy for losses or damages resulting from this product (including claims based on contract, negligence, strict liability, or other legal theories), shall be limited to, at Seller's election, one of the following:

1. Refund of purchase price paid by buyer or user for product bought, or
2. Replacement of amount of product used.

Seller shall not be liable for losses or damages resulting from handling or use of this product unless Seller is promptly notified of such loss or damage in writing. In no case shall Seller be liable for consequential or incidental damages or losses.

The terms of the Warranty Disclaimer and Inherent Risks of Use above and this Limitation of Remedies cannot be varied by any written or verbal statements or agreements. No employee or sales agent of Seller or the Seller is authorized to vary or exceed the terms of the Warranty Disclaimer or this Limitation of Remedies in any manner.

**DISTRIBUTED BY:**  
**LEBANON SEABOARD CORPORATION**  
 1600 EAST CUMBERLAND STREET  
 LEBANON, PA 17042  
[www.LebanonTurf.com](http://www.LebanonTurf.com)



**39201**

SPECIMEN LABEL